

# New Hypothesis

## Kvarky

<b><i>b</i></b>	<b><i>t</i></b>	<b><i>u</i></b>	<b><i>d</i></b>	<b><i>s</i></b>	<b><i>c</i></b>
$\frac{x^3 \cdot t^{5/3}}{x^2 \cdot t^{7/3}}$	$\frac{x^3 \cdot t^{8/3}}{x^2 \cdot t^{10/3}}$	$\frac{x^1 \cdot t^{-1/3}}{x^0 \cdot t^{+1/3}}$	$\frac{x^1 \cdot t^{2/3}}{x^0 \cdot t^{4/3}}$	$\frac{x^2 \cdot t^{2/3}}{x^1 \cdot t^{4/3}}$	$\frac{x^2 \cdot t^{5/3}}{x^1 \cdot t^{7/3}}$

## Leptony

## Antileptony

<p>(e<sup>-</sup>)</p> $\frac{t^1 \cdot x^2 \cdot t^1}{1 \cdot x^2 \cdot t^1} = \frac{x^2 \cdot t^2}{x^2 \cdot t^1}$ <p>(τ<sup>-</sup>)</p> $\frac{t^1 \cdot x^2 \cdot t^0}{1 \cdot x^2 \cdot t^0} = \frac{x^2 \cdot t^1}{x^2 \cdot t^0}$ <p>(μ<sup>-</sup>)</p> $\frac{t^1 \cdot x^1 \cdot t^1}{1 \cdot x^1 \cdot t^1} = \frac{x^1 \cdot t^2}{x^1 \cdot t^1}$ <p>(ν<sub>μ</sub>)<sup>0</sup></p> $\frac{t^1 \cdot x^1 \cdot t^0}{1 \cdot x^1 \cdot t^0} = \frac{x^1 \cdot t^1}{x^1 \cdot t^0}$ <p>(ν<sub>τ</sub>)<sup>0</sup></p> $\frac{t^1 \cdot x^0 \cdot t^1}{1 \cdot x^0 \cdot t^1} = \frac{x^0 \cdot t^2}{x^0 \cdot t^1}$ <p>(ν<sub>e</sub>)<sup>0</sup></p> $\frac{t^1 \cdot x^0 \cdot t^0}{1 \cdot x^0 \cdot t^0} = \frac{x^0 \cdot t^1}{x^0 \cdot t^0}$	<p>(e<sup>+</sup>)</p> $\frac{1 \cdot x^2 \cdot t^1}{t^1 \cdot x^2 \cdot t^1} = \frac{x^2 \cdot t^1}{x^2 \cdot t^2}$ <p>(τ<sup>+</sup>)</p> $\frac{1 \cdot x^2 \cdot t^0}{t^1 \cdot x^2 \cdot t^0} = \frac{x^2 \cdot t^0}{x^2 \cdot t^1}$ <p>(μ<sup>+</sup>)</p> $\frac{1 \cdot x^1 \cdot t^1}{t^1 \cdot x^1 \cdot t^1} = \frac{x^1 \cdot t^1}{x^1 \cdot t^2}$ <p>(ν<sub>μ</sub><sup>~</sup>)<sup>0</sup></p> $\frac{1 \cdot x^1 \cdot t^0}{t^1 \cdot x^1 \cdot t^0} = \frac{x^1 \cdot t^0}{x^1 \cdot t^1}$ <p>(ν<sub>τ</sub><sup>~</sup>)<sup>0</sup></p> $\frac{1 \cdot x^0 \cdot t^1}{t^1 \cdot x^0 \cdot t^1} = \frac{x^0 \cdot t^1}{x^0 \cdot t^2}$ <p>(ν<sub>e</sub><sup>~</sup>)<sup>0</sup></p> $\frac{1 \cdot x^0 \cdot t^0}{t^1 \cdot x^0 \cdot t^0} = \frac{x^0 \cdot t^0}{x^0 \cdot t^1}$
---	---